

MAY FREELY DISTRIBUTE PRINTED HANDOUTS. FULL DETAILS AT http://Skymaps.com/terms.html

## About the Celestial Objects

Listed on this page are several of the brighter, more interesting celestial objects visible in the evening sky this month (refer to the monthly sky map). The objects are grouped into three categories. Those that can be easily seen with the naked eye (that is, without optical aid), those easily seen with binoculars, and those requiring a telescope to be appreciated. Note, all of the objects (except single stars) will appear more impressive when viewed through a telescope or very large **binoculars.** They are grouped in this way to highlight objects that can be seen using the optical equipment that may be available to the star gazer.

## Tips for Observing the Night Sky

When observing the night sky, and in particular deep-sky objects such as star clusters, nebulae, and galaxies, it's always best to observe from a dark location. Avoid direct light from street lights and other sources. If possible observe from a dark location away from the light pollution that surrounds many of today's large cities.

You will see more stars after your eyes adapt to the darkness—usually about 10 to 20 minutes after you go outside. Also, if you need to use a torch to view the sky map, cover the light bulb with red cellophane. This will preserve your dark vision.

Finally, even though the Moon is one of the most stunning objects to view through a telescope, its light is so bright that it brightens the sky and makes many of the fainter objects very difficult to see. So try to observe the evening sky on moonless nights around either New Moon or Last Quarter.

## Astronomical Glossarv

**Conjunction** – An alignment of two celestial bodies such that they present the least angular separation as viewed from Earth.

**Constellation** – A defined area of the sky containing a star pattern.

**Diffuse Nebula** – A cloud of gas illuminated by nearby stars.

**Double Star** – Two stars that appear close to each other in the sky; either linked by gravity so that they orbit each other (binary star) or lying at different distances from Earth (optical double). Apparent separation of stars is given in seconds of arc (").

Ecliptic – The path of the Sun's center on the celestial sphere as seen from Earth.

**Elongation** – The angular separation of two celestial bodies. For Mercury and Venus the greatest elongation occurs when they are at their most angular distance from the Sun as viewed from Earth.

**Galaxy** – A mass of up to several billion stars held together by gravity.

Globular Star Cluster – A ball-shaped group of several thousand old stars.

Light Year (ly) - The distance a beam of light travels at 300,000 km/sec in one vear.

Magnitude – The brightness of a celestial object as it appears in the sky.

**Open Star Cluster** – A group of tens or hundreds of relatively young stars.

**Opposition** – When a celestial body is opposite the Sun in the sky.

**Planetary Nebula** – The remnants of a shell of gas blown off by a star.

Universal Time (UT) - A time system used by astronomers. Also known as Greenwich Mean Time. USA Eastern Standard Time (for example, New York) is 5 hours behind UT.

## **0CT0BER 2023** Easily Seen with the Naked Eve

**NORTHERN HEMISPHERE** 

CIS

**OBJE** 

LESTIAL

Sol

ma

13

~

☆

Easily O		with the Nuneu Lyc
Altair Capella Arcturus δ Cephei Deneb α Herculis Vega Algol Fomalhaut Pleiades Polaris	Aql Aur Boo Cep Cyg Her Lyr Per PsA Tau UMi	<ul> <li>Brightest star in Aquila. Name means "the flying eagle". Dist=16.7 ly.</li> <li>The 6th brightest star. Appears yellowish in color. Spectroscopic binary. Dist=42 ly.</li> <li>Orange, giant K star. Name means "bear watcher". Dist=36.7 ly.</li> <li>Cepheid prototype. Mag varies between 3.5 &amp; 4.4 over 5.366 days. Mag 6 companion.</li> <li>Brightest star in Cygnus. One of the greatest known supergiants. Dist=1,400±200 ly.</li> <li>Semi-regular variable. Magnitude varies between 3.1 &amp; 3.9 over 90 days. Mag 5.4 companion.</li> <li>The 5th brightest star in the sky. A blue-white star. Dist=25.0 ly.</li> <li>Famous eclipsing binary star. Magnitude varies between 2.1 &amp; 3.4 over 2.867 days.</li> <li>Brightest star in Piscis Austrinus. In Arabic the "fish's mouth". Dist=25 ly.</li> <li>The Seven Sisters. Spectacular cluster. Many more stars visible in binoculars. Dist=380 ly.</li> <li>The North Pole Star. A telescope reveals an unrelated mag 8 companion star. Dist=433 ly.</li> </ul>
Easily Se	een	with Binoculars
M8 M25 M22 Mizar & Alcor	Sgr Sgr Sgr UMa	<ul> <li>The Andromeda Galaxy. Most distant object visible to naked eye. Dist=2.93 million ly.</li> <li>Resembles a fuzzy star in binoculars.</li> <li>Bright Cepheid variable. Mag varies between 3.6 &amp; 4.5 over 7.166 days. Dist=1,200 ly.</li> <li>Herschel's Garnet Star. One of the reddest stars. Mag 3.4 to 5.1 over 730 days.</li> <li>Long period pulsating red giant. Magnitude varies between 3.3 &amp; 14.2 over 407 days.</li> <li>May be visible to the naked eye under good conditions. Dist=900 ly.</li> <li>Wide pair of white stars. One of the finest binocular pairs in the sky. Dist=100 ly.</li> <li>Best globular in northern skies. Discovered by Halley in 1714. Dist=23,000 ly.</li> <li>Fainter and smaller than M13. Use a telescope to resolve its stars.</li> <li>Famous Double Double. Binoculars show a double star. High power reveals each a double.</li> <li>Semi-regular variable. Magnitude varies between 3.9 &amp; 5.0 over 46.0 days.</li> <li>Close to the brighter M10. Dist=18,000 ly.</li> <li>3 degrees from the fainter M12. Both may be glimpsed in binoculars. Dist=14,000 ly.</li> <li>Large, scattered open cluster. Visible with binoculars.</li> <li>Scattered open cluster. Visible with binoculars.</li> <li>Moly globular known to contain a planetary nebula (Mag 14, d=1"). Dist=30,000 ly.</li> <li>Double Cluster in Perseus. NGC 869 &amp; 884. Excellent in binoculars. Dist=7,300 ly.</li> <li>Bright cluster located about 6 deg N of "teapot's" lid. Dist=1,900 ly.</li> <li>A spectacular globular star cluster. Telescope will show stars. Dist=10,000 ly.</li> <li>Good eyesight or binoculars reveals 2 stars. Not a binary. Mizar has a mag 4 companion.</li> </ul>
		© Coathanger asterism or "Brocchi's Cluster". Not a true star cluster. Dist=218 to 1,140 ly.
$\begin{array}{l} \gamma \text{ Andromedae} \\ 7009 \\ 7293 \\ \gamma \text{ Arietis} \\ \text{M51} \\ \eta \text{ Cassiopeiae} \\ \text{Albireo} \\ \text{61 Cygni} \\ \gamma \text{ Delphini} \\ \beta \text{ Lyrae} \\ \text{M57} \\ \text{M23} \\ \text{M20} \\ \text{M21} \\ \text{M11} \\ \text{M16} \\ \text{M33} \\ \text{M81} \\ \text{M27} \end{array}$	Aqr Aqr CVn Cas Cyg Del Lyr Lyr Sgr Sgr Sgr Sgr Sct Ser Tri	<ul> <li>Attractive double star. Bright orange star with mag 5 blue companion. Sep=9.8".</li> <li>Saturn Nebula. Requires 8-inch telescope to see Saturn-like appendages.</li> <li>Helix Nebula. Spans nearly 1/4 deg. Requires dark sky. Dist=300 ly.</li> <li>Impressive looking double blue-white star. Visible in a small telescope. Sep=7.8".</li> <li>Whirlpool Galaxy. First recognised to have spiral structure. Dist=25 million ly.</li> <li>Yellow star mag 3.4 &amp; orange star mag 7.5. Dist=19 ly. Orbit=480 years. Sep=12".</li> <li>Beautiful double star. Contrasting colours of orange and blue-green. Sep=34.4".</li> <li>Attractive double star. Mags 5.2 &amp; 6.1 orange dwarfs. Dist=11.4 ly. Sep=28.4".</li> <li>Appear yellow &amp; white. Mags 4.3 &amp; 5.2. Dist=100 ly. Struce 2725 double in same field.</li> <li>Eclipsing binary. Mag varies between 3.3 &amp; 4.3 over 12.940 days. Fainter mag 7.2 blue star.</li> <li>Ring Nebula. Magnificent object. Smoke-ring shape. Dist=4,100 ly.</li> <li>Elongated star cluster. Telescope required to show stars. Dist=2,100 ly.</li> <li>Trifid Nebula. A telescope shows 3 dust lanes trisecting nebula. Dist=5,200 ly.</li> <li>A fine and impressive cluster. Dist=4,200 ly.</li> <li>Omega Nebula. Contains the star cluster NGC 6618. Dist=4,900 ly.</li> <li>Wild Duck Cluster. Resembles a globular through binoculars. V-shaped. Dist=5,600 ly.</li> <li>Eagle Nebula. Requires a telescope of large aperture. Dist=8,150 ly.</li> <li>Fine face-on spiral galaxy. Requires a large aperture telescope. Dist=2.3 million ly.</li> <li>Beautiful spiral galaxy visible with binoculars. Easy to see in a telescope.</li> <li>Dumbbell Nebula. Large, twin-lobed shape. Most spectacular planetary. Dist=975 ly.</li> </ul>
	Altair Capella Arcturus $\delta$ Cephei Deneb $\alpha$ Herculis Vega Algol Fomalhaut Pleiades Polaris <b>Easily So</b> M31 M2 $\eta$ Aquilae $\mu$ Cephei $\chi$ Cygni M39 $\nu$ Draconis M13 M92 $\epsilon$ Lyrae M12 M10 IC 4665 6633 M15 Double Cluster M8 M25 M22 Mizar & Alcor Cr 399 <b>Telescoj</b> $\gamma$ Andromedae 7009 7293 $\gamma$ Arietis M51 $\eta$ Cassiopeiae Albireo 61 Cygni $\gamma$ Delphini $\beta$ Lyrae M57 M23 M21 M17 M11 M16 M33 M81	AltairAqlCapellaAurArcturusBooδ CepheiCepDenebCyg $\alpha$ HerculisHerVegaLyrAlgolPerFomalhautPsAPleiadesTauPolarisUMi <b>Easily Seen</b> M31AndM2Aqr $\eta$ AquilaeAql $\mu$ CepheiCep $\chi$ CygniCygM39Cyg $\gamma$ DraconisDraM13HerM92Her $klyrae$ LyrM12OphM15PegDouble ClusterPerM8SgrM22SgrM13AqrY AndromedaeAqrY ArietisAriM51CVn $\gamma$ CassiopeiaeCasAlbireoCygG1 CygniCyg $\beta$ LyraeLyrM51CVn $\gamma$ CassiopeiaeCasAlbireoCygG1 CygniCyg $\beta$ LyraeLyrM51Cyn $\gamma$ DelphiniDel $\beta$ LyraeLyrM51SgrM23SgrM21SgrM21SgrM33TriM81UMa

The Evening Sky Map (ISSN 1839-7735) Copyright © 2000-2023 Kym Thalassoudis. All Rights Reserved.